

(19)

Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 955 293 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:

10.11.1999 Bulletin 1999/45

(51) Int. Cl.⁶: C07D 231/40

(21) Application number: 97946071.4

(86) International application number:
PCT/JP97/04399

(22) Date of filing: 02.12.1997

(87) International publication number:
WO 98/24768 (11.06.1998 Gazette 1998/23)(84) Designated Contracting States:
DE FR GB IT

(30) Priority: 03.12.1996 JP 33759396

(71) Applicant:

BANYU PHARMACEUTICAL CO., LTD.
Chuo-ku, Tokyo 103-8416 (JP)

- KANATANI, Akio,
Banyu Pharmaceutical Co., Ltd
Tsukuba-shi, Ibaraki 300-26 (JP)
- IHARA, Masaki,
Banyu Pharmaceutical Co., Ltd
Tsukuba-shi, Ibaraki 300-26 (JP)

(72) Inventors:


- FUKAMI, Takehiro,
Banyu Pharmaceutical Co., Ltd
Tsukuba-shi, Ibaraki 300-26 (JP)
- FUKURODA, Takahiro,
Banyu Pharmaceutical Co., Ltd
Tsukuba-shi, Ibaraki 300-26 (JP)

(74) Representative:

Woods, Geoffrey Corlett et al
J.A. KEMP & CO.
Gray's Inn
14 South Square
London WC1R 5LX (GB)

(54) NOVEL UREA DERIVATIVES

(57) The present invention relates to a compound represented by the general formula [I]:



wherein A represents a nitrogen atom or a group represented by C-R⁵; Ar¹ represents an aryl group which may have a substituent selected from the group consisting of a halogen atom and lower alkyl and lower haloalkyl groups; Ar² represents an aryl or heteroaryl group which may have a substituent selected from the group consisting of a halogen atom and lower alkyl, lower alkenyl, lower haloalkyl, lower alkoxy, lower alkylthio, lower alkylamino, lower dialkylamino and aryl groups; R¹ represents a hydrogen atom, a lower alkyl group or a bond formed by linking to R⁵; R² represents a hydrogen atom or a lower alkyl group; R³ and R⁴ may be the same or different and each represents a hydrogen atom or a lower alkyl group; and R⁵ represents a hydrogen atom or a hydroxyl, lower alkyl or lower alkoxy group or a bond formed by linking to R¹, or its salt, a process for its preparation and an agent for the treatment of bulimia, obesity or diabetes comprising it as an active ingredient.

EP 0 955 293 A1

Description

FIELD OF THE INVENTION

- 5 [0001] The present invention is useful in the field of medicines. More specifically, novel pyrazole derivatives of the present invention are useful as neuropeptide Y receptor antagonists and as agents for the treatment of various diseases of circulatory organs, central nervous system and metabolic system.

BACKGROUND OF THE INVENTION

- 10 [0002] Neuropeptide Y (to be referred to as NPY hereinafter) is a peptide consisting of 36 amino acids, which was isolated from porcine brain for the first time by Tatemoto et al. in 1982 [Nature, vol.296, p.659 (1982)]. NPY is broadly distributed in central nervous and peripheral nervous systems and has various *in vivo* functions as one of the peptides most abundantly present in the nervous system. That is, in the central nervous system, NPY acts as an aperitive and significantly promotes a fat accumulation via secretion of various hormones and actions of the nervous system. It is known that a continuous intracerebroventricular administration of NPY induces obesity and insulin resistance based on the above actions. NPY is also associated with the control of mood and functions of the central autonomic nervous system. In addition, in the peripheral nervous system, NPY is present together with norepinephrine in the sympathetic nerve terminal and associated with the tension of the sympathetic nervous system. It is known that a peripheral administration of NPY causes vasoconstriction and enhances actions of other vasoconstrictors including norepinephrine [International Journal of Obesity, vol.19, p.517 (1995); Endocrinology, vol.133, p.1753 (1993); British Journal of Pharmacology, vol.95, p.419 (1988)].

- 25 [0003] The function of NPY is expressed when it is bound to an NPY receptor present in the central or peripheral nervous system. Therefore, the expression of the function of NPY can be prevented if the binding of NPY to the NPY receptor is inhibited. Consequently, it is expected that compounds capable of inhibiting the binding of NPY to the NPY receptor are useful in the prevention or treatment of various diseases associated with NPY, for example, diseases of circulatory organs such as hypertension, nephropathy, cardiopathy and angiospasm; diseases of central nervous system such as bulimia, depression, epilepsy and dementia; metabolic diseases such as obesity, diabetes and dysendocrisis, or glaucoma [Trends in Pharmacological Sciences, vol.15, p.153 (1994)].


- 30 [0004] Compounds structurally similar to the compounds of the present invention are disclosed in International Publication WO 96/14843, JP 3093774A, JP 2300173A, JP 51146465A and etc. However, these publications do not clearly disclose nor suggest the compound of the present invention. And, an antagonistic action to NPY is not described at all therein.

DISCLOSURE OF THE INVENTION

- 35 [0005] An object of the present invention is to provide a new medicine having an antagonistic action to NPY.

- [0006] The present inventors have found that a compound represented by the general formula [I]:

40



45

- 50 wherein A represents a nitrogen atom or a group represented by C-R⁵; Ar¹ represents an aryl group which may have a substituent selected from the group consisting of a halogen atom and lower alkyl and lower haloalkyl groups; Ar² represents an aryl or heteroaryl group which may have a substituent selected from the group consisting of a halogen atom and lower alkyl, lower alkenyl, lower haloalkyl, lower alkoxy, lower alkylthio, lower alkylamino, lower dialkylamino and aryl groups; R¹ represents a hydrogen atom, a lower alkyl group or a bond formed by linking to R⁵; R² represents a hydrogen atom or a lower alkyl group; R³ and R⁴ may be the same or different and each represents a hydrogen atom or a lower alkyl group; or R³ and R⁴ are linked to each other to form an alkylene group containing 2 to 4 carbon atoms which may have a lower alkyl group; and R⁵ represents a hydrogen atom or a hydroxyl, lower alkyl or lower alkoxy group or a bond formed by linking to R¹, has an antagonistic action to NPY.

[0007] Since the compound [I] of the present invention has an antagonistic action to NPY, it is useful as an agent for the treatment of various diseases associated with NPY, for example, diseases of circulatory organs such as hypertension, nephropathy, cardiopathy and angiospasm, diseases of central nervous system such as bulimia, depression, epilepsy and dementia, metabolic diseases such as obesity, diabetes and dysendocrisiasis, or glaucoma.

5 [0008] Especially, the compound [I] of the present invention is useful as an agent for the treatment of bulimia, obesity, diabetes or the like.

[0009] The present invention relates to a compound represented by the general formula [I] or its salt as well as a method for its preparation and its use.

[0010] Symbols and terms as used herein are described below.

10 [0011] The term "halogen atom" as used herein means fluorine, chlorine, bromine or iodine.

[0012] The term "lower alkyl group" as used herein means a straight, branched or cyclic alkyl group containing 1 to 7 carbon atoms, and its illustrative examples include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, tert-pentyl, 1-methylbutyl, 2-methylbutyl, 1,2-dimethylpropyl, 1-ethylpropyl, hexyl, isohexyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 2,2-dimethylbutyl, 1-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-2-methylpropyl, 1-ethyl-1-methylpropyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclopropylmethyl, 1-cyclopropylethyl, 2-cyclopropylethyl, 1-cyclopropylpropyl, 2-cyclopropylpropyl, 3-cyclopropylpropyl, cyclopentylmethyl, 2-cyclopentylethyl and cyclohexylmethyl and the like.

15 [0013] The term "lower haloalkyl group" as used herein means the above lower alkyl group comprising the aforementioned halogen atom, and its illustrative examples include fluoromethyl, difluoromethyl, trifluoromethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, chloromethyl, dichloromethyl, trichloromethyl, 1-chloroethyl and 2-chloroethyl and the like.

20 [0014] The term "aryl group" as used herein means phenyl, naphthyl or anthryl, of which phenyl and naphthyl are preferred.

[0015] The term "lower alkenyl group" as used herein means a straight or branched alkenyl group containing 2 to 7 carbon atoms, and its illustrative examples include vinyl, 2-propenyl, isopropenyl, 3-butenyl, 2-butenyl, 1-butenyl, 1-methyl-2-propenyl, 1-methyl-1-propenyl, 1-ethyl-1-ethenyl, 2-methyl-2-propenyl, 2-methyl-1-propenyl, 3-methyl-2-butenyl and 4-pentenyl and the like.

30 [0016] The term "lower alkoxy group" as used herein means an alkoxy group comprising the aforementioned lower alkyl group, namely an alkoxy group containing 1 to 7 carbon atoms, or an alkylenedioxy group containing 1 to 3 carbon atoms, and its illustrative examples include methoxy, ethoxy, propyloxy, isopropyloxy, butoxy, isobutyloxy, tert-butoxy, pentyloxy, cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy, cycloheptyloxy, cyclopropylmethyloxy, 1-cyclopropylethyloxy, 2-cyclopropylethyloxy, 1-cyclopropylpropyloxy, 2-cyclopropylpropyloxy, 3-cyclopropylpropyloxy, cyclopentylmethyloxy, 2-cyclopentylethyloxy, cyclohexylmethyloxy, methylenedioxy, ethylenedioxy and trimethylenedioxy and the like.

35 [0017] The term "lower alkylthio group" as used herein means an alkylthio group comprising the aforementioned lower alkyl group, namely an alkylthio group containing 1 to 7 carbon atoms, and its illustrative examples include methylthio, ethylthio, propylthio, isopropylthio, butylthio, isobutylthio, tert-butylthio, pentythio, cyclopropylthio, cyclobutylthio, cyclopentylthio, cyclohexylthio, cycloheptylthio, cyclopropylmethylthio, 1-cyclopropylethylthio, 2-cyclopropylethylthio, 1-cyclopropylpropylthio, 2-cyclopropylpropylthio, 3-cyclopropylpropylthio, cyclopentylmethylthio, 2-cyclopentylethylthio and cyclohexylmethylthio and the like.

[0018] The term "lower alkylamino group" as used herein means an amino group monosubstituted with the aforementioned lower alkyl group, and its illustrative examples include methylamino, ethylamino, propylamino, isopropylamino, butylamino, sec-butylamino and tert-butylamino and the like.

45 [0019] The term "lower dialkylamino group" as used herein means an amino group disubstituted with the aforementioned lower alkyl group, and its illustrative examples include dimethylamino, diethylamino, ethylmethylamino, dipropylamino, methylpropylamino and diisopropylamino and the like.

[0020] The term "heteroaryl group" as used herein means a 5- or 6-membered monocyclic aromatic heterocyclic ring group containing 1 or more, preferably 1 to 3, of heteroatoms which may be the same or different and are selected from the group consisting of oxygen, nitrogen and sulfur, or a condensed aromatic heterocyclic ring group in which said 50 monocyclic aromatic heterocyclic ring group and the aforementioned aryl group are condensed or in which the same or different members of said monocyclic aromatic heterocyclic ring groups are mutually condensed, and its illustrative examples include pyrrolyl, furyl, thienyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, triazolyl, oxadiazolyl, thiadiazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, indazolyl, purinyl, quinolyl, isoquinolyl, phthalazinyl, naphthylidiny, quinoxaliny, quinazolinyl, cinnolinyl and pteridinyl and the like.

55 [0021] The term "alkylene group containing 2 to 4 carbon atoms" as used herein means a straight chain alkylene group, namely ethylene, trimethylene or tetramethylene.

[0022] The "salt" of the compound represented by the general formula [I] means any conventional pharmaceutically

acceptable salt, and its examples include acid addition salts based on basic groups such as a basic heterocyclic ring group or an amino substituents and the like.

[0023] Illustrative examples of the acid addition salt include an inorganic salt such as hydrochloride, sulfate, nitrate, phosphate and perchlorate; an organic salt such as maleate, fumarate, tartarate, citrate, ascorbate and trifluoroacetate; and sulfonate such as methanesulfonate, isethionate, benzenesulfonate and p-toluenesulfonate.

[0024] The term "agent for the treatment" as used herein means a drug to be used for the treatment and/or prevention of various diseases.

[0025] Various symbols in the general formula [I] are described further in detail with reference to its preferred embodiment in order to explain the compound represented by the above general formula [I] of the present invention more clearly.

[0026] "A" means nitrogen atom or a group represented by C-R⁵.

[0027] Nitrogen atom is preferred as "A".

[0028] Ar¹ means an aryl group which may have a substituent selected from the group consisting of a halogen atom and lower alkyl and lower haloalkyl groups.

[0029] The term "an aryl group which may have a substituent selected from the group consisting of a halogen atom and lower alkyl and lower haloalkyl groups" means the aforementioned aryl group unsubstituted or substituted at any substitutable position with 1 or more, preferably 1 or 2, substituents which may be the same or different and are selected from the group consisting of a halogen atom and lower alkyl and lower haloalkyl groups.

[0030] Preferred examples of the halogen atom as said substituent include chlorine and bromine.

[0031] Preferred examples of the lower alkyl group as said substituent include methyl, ethyl, propyl, isopropyl, cyclopropyl, butyl, isobutyl, sec-butyl, tert-butyl and pentyl, of which methyl, ethyl, propyl, isopropyl and cyclopropyl are more preferred.

[0032] Preferred examples of the lower haloalkyl group as said substituent include trifluoromethyl.

[0033] The said substituent is preferably a halogen atom.

[0034] Preferred examples of the aryl group as Ar¹ include phenyl.

[0035] In consequence, preferred examples of Ar¹ include phenyl and 3-chlorophenyl, of which 3-chlorophenyl is more preferred.

[0036] Ar² means an aryl or heteroaryl group which may have a substituent selected from the group consisting of a halogen atom and lower alkyl, lower alkenyl, lower haloalkyl, lower alkoxy, lower alkylthio, lower alkylamino, lower dialkylamino and aryl groups.

[0037] The term "an aryl or heteroaryl group which may have a substituent selected from the group consisting of a halogen atom and lower alkyl, lower alkenyl, lower haloalkyl, lower alkoxy, lower alkylthio, lower alkylamino, lower dialkylamino and aryl groups" means the aforementioned aryl or heteroaryl group unsubstituted or substituted at any substitutable position with 1 or more, preferably 1 or 2, substituents which may be the same or different and are selected from the group consisting of a halogen atom and lower alkyl, lower alkenyl, lower haloalkyl, lower alkoxy, lower alkylthio, lower alkylamino, lower dialkylamino and aryl groups.

[0038] Preferred examples of the halogen atom as said substituent include chlorine and bromine.

[0039] Preferred examples of the lower alkyl group as said substituent include methyl and ethyl.

[0040] Preferred examples of the lower alkenyl group as said substituent include vinyl, 2-propenyl, isopropenyl, 2-butenyl and 3-methyl-2-butenyl, of which vinyl and 2-propenyl are more preferred.

[0041] Preferred examples of the lower haloalkyl group as said substituent include fluoromethyl and trifluoromethyl.

[0042] Preferred examples of the lower alkoxy group as said substituent include methoxy, ethoxy, propyloxy and methylenedioxy, of which methoxy and methylenedioxy are more preferred.

[0043] Preferred examples of the lower alkylthio group as said substituent include methylthio, ethylthio and propylthio, of which methylthio is more preferred.

[0044] Preferred examples of the lower alkylamino group as said substituent include methylamino.

[0045] Preferred examples of the lower dialkylamino group as said substituent include dimethylamino.

[0046] Preferred examples of the aryl group as said substituent include phenyl.

[0047] As said substituent, a halogen atom and a lower alkyl, lower alkoxy, lower alkylthio or lower dialkylamino group are preferred.

[0048] Preferred examples of the aryl group as Ar² include phenyl, and those of the heteroaryl group include pyridyl.

[0049] In consequence, preferred examples of Ar² include phenyl, 1-naphthyl, 2-naphthyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 3,4-dichlorophenyl, 2-bromophenyl, 3-bromophenyl, 4-bromophenyl, 5-chloro-3-pyridyl, 5-bromo-3-pyridyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 3-ethylphenyl, 4-ethylphenyl, 4-propylphenyl, 5-methyl-2-pyridyl, 6-methyl-3-pyridyl, 6-ethyl-3-pyridyl, 2-methyl-4-pyridyl, 2-ethyl-4-pyridyl, 2-propyl-4-pyridyl, 2-butyl-4-pyridyl, 3-methyl-4-pyridyl, 3-vinylphenyl, 4-vinylphenyl, 4-(2-propenyl)phenyl, 5-vinyl-3-pyridyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 4-ethoxyphenyl, 4-isopropoxyphenyl, 3,4-dimethoxyphenyl, 3,4-methylenedioxyphenyl, 6-methoxy-3-pyridyl, 2-meth-

oxy-4-pyridyl, 3-dimethylaminophenyl, 4-dimethylaminophenyl, 4-methoxy-3-dimethylaminophenyl, 2-methylthiophenyl, 3-methylthiophenyl, 4-methylthiophenyl, 2-methylthio-4-pyridyl, 3-biphenyl and 4-biphenyl, of which 4-pyridyl, 2-methyl-4-pyridyl, 2-ethyl-4-pyridyl, 2-propyl-4-pyridyl, 2-butyl-4-pyridyl, 3-chlorophenyl, 4-chlorophenyl, 3,4-dichlorophenyl, 3-bromophenyl, 4-bromophenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 4-ethylphenyl, 3-methyl-4-pyridyl, 3-ethyl-4-pyridyl, 4-vinylphenyl, 4-(2-propenyl)phenyl, 3-methoxyphenyl, 4-methoxyphenyl, 4-ethoxyphenyl, 3,4-dimethoxyphenyl, 3,4-methylenedioxyphenyl, 3-methoxy-4-pyridyl, 4-methylthiophenyl and 3-methylthio-4-pyridyl are more preferred.

[0050] R¹ means a hydrogen atom, a lower alkyl group or a bond formed by linking to R⁵.

[0051] Preferred examples of the lower alkyl group as R¹ include methyl and ethyl.

[0052] The term "a bond formed by linking to R⁵" means that, when A is a group represented by C-R⁵, R¹ and R⁵ are linked to each other to form a double bond together with the existing bond.

[0053] Preferred as R¹ is hydrogen or a bond formed by linking to R⁵.

[0054] R² means a hydrogen atom or a lower alkyl group.

[0055] Preferred examples of the lower alkyl group as R² include methyl and ethyl.

[0056] Hydrogen is preferred as R².

[0057] R³ and R⁴ may be the same or different and each means a hydrogen atom or a lower alkyl group, or R³ and R⁴ are linked to each other to form an alkylene group containing 2 to 4 carbon atoms which may have a lower alkyl group.

[0058] Preferred examples of the lower alkyl group as R³ and R⁴ include methyl and ethyl.

[0059] The term "an alkylene group containing 2 to 4 carbon atoms which may have a lower alkyl group" means the aforementioned alkylene group unsubstituted or substituted at any substitutable position with 1 or more of the aforementioned lower alkyl groups which may be the same or different, of which the unsubstituted alkylene group is preferred.

[0060] Preferred examples of the lower alkyl group as the substituent include methyl, ethyl, propyl and isopropyl, of which methyl is more preferred.

[0061] Preferred examples of R³ and R⁴ include a lower alkyl group which may be the same or different, and an alkylene group containing 2 to 4, particularly 2, carbon atoms which is formed by linking R³ and R⁴ and which may have a lower alkyl group.

[0062] R⁵ means a hydrogen atom, a hydroxyl, lower alkyl or lower alkoxy group or a bond formed by linking to R¹.

[0063] Preferred examples of the lower alkyl group as R⁵ include methyl and ethyl.


[0064] Preferred examples of the lower alkoxy group as R⁵ include methoxy, ethoxy and propyloxy.

[0065] The term "a bond formed by linking to R¹" means that, when A is a group represented by C-R⁵, R¹ and R⁵ are linked to each other to form a double bond together with the existing bond.

[0066] Preferred as R⁵ is hydrogen, hydroxyl or a bond formed by linking to R¹.

[0067] Depending on the nature of substituent, the compounds of the present invention may exist in various stereoisomers including optical isomers, diastereomers and geometrical isomers, and tautomers. All of these stereoisomers and tautomers and their mixtures are also included in the present invention.

[0068] In this connection, in order to avoid unnecessary confusion in naming each of the compounds of the present invention, position numbers of the pyrazole ring moiety of the compound represented by the general formula [I] are defined as shown in the following general formula [I'], and the nomenclature and other explanations of each compound are described based on this formula.



[0069] Illustrative examples of the compound represented by the general formula [I] include:

5-(4-methoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-chlorophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(2-methylphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,


5-(3-methylphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-methylphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(2-methoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(3,4-dichlorophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-bromophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(3-chlorophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-phenyl-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-dimethylaminophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(3-dimethylaminophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(3,4-dimethoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-isopropoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-ethoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(3-bromophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 3-[4-(3-chlorophenyl)piperazinyl]carbonylamino-5-(4-methoxyphenyl)pyrazole,
 3-[4-(3-chlorophenyl)piperazinyl]carbonylamino-5-(3,4-dimethoxyphenyl)pyrazole,
 5-(2-chlorophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-methylthiophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(3,4-methylenedioxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 3-(4-phenylpiperazinyl)carbonylamino-5-(4-pyridyl)pyrazole,
 5-(4-methoxyphenyl)-3-methyl[2-(methylphenylamino)ethyl]aminocarbonylaminopyrazole,
 5-(3-methoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-methoxyphenyl)-3-(2-methyl-4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-methoxyphenyl)-3-[2-(phenylamino)ethyl]aminocarbonylaminopyrazole,
 5-(4-biphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(3-dimethylamino-4-methoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 3-(4-hydroxy-4-phenylpiperidino)carbonylamino-5-(4-methoxyphenyl)pyrazole,
 5-(4-methoxyphenyl)-3-(4-phenyl-1,2,3,6-tetrahydropyridin-1-yl)carbonylaminopyrazole,
 5-(4-methoxyphenyl)-3-(4-phenylpiperidino)carbonylaminopyrazole,
 5-(4-methoxyphenyl)-3-(4-phenylhomopiperazinyl)carbonylaminopyrazole.

[0070] Next, a process for the preparation of the compound of the present invention is described below.

[0071] The compound [I] of the present invention can be prepared, for example, by the following process or the method as shown in Examples. However, the process for the preparation of the compound [I] of the present invention is not limited thereto.

Process


[0072] The compound represented by the general formula [I] can be prepared by reacting a compound represented by the general formula [II]:



[II]

wherein Ar² is as defined above,


with a compound represented by the general formula [III]:




[III]

wherein Ar³ represents a phenyl group which may be substituted with a halogen atom or a nitro group; and X represents

a halogen atom, to obtain a compound represented by the general formula [IV]:



10 wherein Ar^2 and Ar^3 are as defined above,
and then reacting the compound [IV] with a compound represented by the general formula [V]:



wherein A, Ar^1 , R^1 , R^2 , R^3 and R^4 are as defined above.

25 [0073] The reaction of the compound of the general formula [II] with the compound of the general formula [III] is generally carried out using 0.5 mole to excess moles, preferably 1 mole to 1.5 moles, of the compound of the general formula [III] with respect to 1 mole of the compound of the general formula [II].

[0074] The reaction is generally carried out in an inert solvent, and preferred examples of said inert solvent include methylene chloride, chloroform, tetrahydrofuran, ethyl ether, benzene, toluene and dimethylformamide and a mixture thereof.

30 [0075] Also, it is desirable to carry out the reaction in the presence of a base, and examples of said base to be used include an organic base such as triethylamine, diisopropylethylamine, pyridine or 4-dimethylaminopyridine, or an inorganic base such as sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate or sodium bicarbonate.

35 [0076] It is desirable to use said base generally in an amount of from 1 mole to excess moles with respect to 1 mole of the compound of the general formula [II]. When said base is liquid, it can be used as both a solvent and a base.

[0077] The reaction temperature is generally within the range of from -78°C to 100°C , preferably from -20°C to 50°C .

[0078] The reaction time is generally within the range of from 5 minutes to 7 days, preferably from 30 minutes to 24 hours.

40 [0079] The reaction of the compound of the general formula [IV] with the compound of the general formula [V] is generally carried out using 1 mole to excess moles, preferably 1 mole to 1.5 moles, of the compound of the general formula [V] with respect to 1 mole of the compound of the general formula [IV] optionally after the isolation of the compound of the general formula [IV] obtained in the above reaction.

[0080] The reaction is generally carried out in an inert solvent, and preferred examples of said inert solvent include methylene chloride, chloroform, tetrahydrofuran and dimethylformamide and a mixture thereof.

45 [0081] Also, it is desirable to carry out the reaction in the presence of a base, and preferred examples of said base to be used include triethylamine and diisopropylethylamine.

[0082] Said base is generally used in an amount of from 1 mole to excess moles, preferably 1 mole to 5 moles, with respect to 1 mole of the compound of the general formula [IV].

[0083] The reaction temperature is generally within the range of from -30°C to 200°C , preferably from 20°C to 100°C .


50 [0084] The reaction time is generally within the range of from 5 minutes to 7 days, preferably from 30 minutes to 24 hours.

[0085] The objective compound in each step can be easily isolated and purified by any conventional separation methods. Examples of such methods include solvent extraction, recrystallization, column chromatography and preparative thin layer chromatography.

55 [0086] The compound represented by the general formula [I] can be converted into its pharmaceutically acceptable salt according to any conventional method and vice versa.

[0087] The compound of the general formula [II] to be used in the present invention is commercially available, or it can be prepared, for example, in accordance with the methods described in references such as Comprehensive Hete-

rocyclic Chemistry, vol.5, edited by A.R. Katritzky, Pergamon Press (1984) or their modifications, or alternatively with the following process or the method as described in the reference examples.



wherein R represents a lower alkyl group; and Ar^2 and X are as defined above.

[0088] According to this process, the compound of the general formula [II] can be prepared by reacting a compound represented by the general formula (1) with acetonitrile in the presence of any base to obtain a compound represented by the general formula (3), and subsequently reacting said compound (3) with hydrazine.

[0089] Conditions for the reaction of the compound (1) with acetonitrile vary depending on the nature of the base to be used.

[0090] For example, when n-butyllithium, lithium diisopropylamide or the like is used as said base, the reaction is generally carried out in any inert solvent such as tetrahydrofuran or ethyl ether at a temperature of from -78°C to room temperature for a reaction period of from 30 minutes to 6 hours.

[0091] When sodium hydride or the like is used as said base, the reaction is generally carried out in an inert solvent such as tetrahydrofuran, ethyl ether or dimethylformamide at a temperature of from room temperature to 100°C for a reaction period of from 1 hour to 6 hours.

[0092] The reaction of the compound (3) with hydrazine is generally carried out in an inert solvent such as ethanol, propanol, isoamyl alcohol, acetic acid, benzene, toluene or xylene or a mixture thereof, using hydrazine in an amount of from 0.5 mole to 10 moles, preferably from 1 mole to 1.5 moles, with respect to 1 mole of the compound (3).

[0093] The reaction temperature is generally from room temperature to the boiling point of the solvent to be used, preferably from 50°C to the boiling point of the solvent to be used.

[0094] The reaction time is generally from 30 minutes to 7 days, preferably from 1 hour to 48 hours.

[0095] The hydrazine to be used in the reaction may be either anhydride or hydrate.

[0096] In addition, the compound of the general formula (3) can also be prepared by using a compound represented by the general formula (2) in stead of the compound represented by the general formula (1), and reacting it with a cyanide.

[0097] In this connection, the compound represented by the general formula (1) or (2) is commercially available or it can be prepared in accordance with known methods or their modifications which may be suitably combined if necessary.

[0098] The compound represented by the general formula [III] or [V] to be used in the present invention is commercially available or it can be prepared according to known methods or their modifications which may be suitably combined if necessary.

[0099] The usefulness of the compound of the present invention as a medicine is demonstrated by showing its antagonistic activity to NPY in the following pharmacological test examples.

Pharmacological Test Example 1 (test of inhibition of NPY binding)

[0100] cDNA Sequence encoding a human NPY Y5 receptor [International Publication WO 96/16542] was cloned into expression vectors pcDNA3, pRc/RSV (manufactured by Invitrogen) and pCI-neo (manufactured by Promega). Using the cationic lipid method [see Proceedings of the National Academy of Science of the United States of America, vol.84, p.7413 (1987)], host cells COS-7, CHO and LM(tk-) (American Type Culture Collection) were transfected with the thus prepared expression vectors to obtain cells in which the NPY Y5 receptor had been expressed.

[0101] Each of the membrane preparations thus prepared from the cells in which the NPY Y5 receptor had been expressed was incubated together with each compound to be tested and 20,000 cpm of [¹²⁵I] peptide YY (manufactured by Amersham) at 25°C for 2 hours in an assay buffer solution (25 mM HEPES buffer, pH 7.4, containing 10 mM magnesium chloride, 1 mM phenylmethylsulfonyl fluoride and 0.1% bacitracin) and then, the reaction mixture was filtered through a glass filter GF/C. After washing with 50 mM Tris buffer, pH 7.4, containing 0.3% BSA, radioactivity on the glass filter was measured using a gamma counter. Non-specific binding was measured in the presence of 1 μM of peptide YY to calculate a concentration of each compound tested which is needed to inhibit 50% of the specific binding of peptide YY (IC₅₀ value) [see Endocrinology, vol.131, p.2090 (1992)]. As the result, IC₅₀ value of the compound of Example 17 was calculated to be 27 nM.

[0102] As shown in the above, the compound of the present invention strongly inhibited the binding of the peptide YY (a homologue of NPY) to the NPY Y5 receptor.

20 Pharmacological Test Example 2 (test of inhibition of feeding behavior induced by bPP)

[0103] Under pentobarbital anesthesia (single intraperitoneal injection of 50mg/kg), a chronic guide cannula (outer diameter 0.8 mm; inner diameter 0.5 mm; length 10 mm) was stereotactically inserted in a right lateral cerebral ventricle of each of SD male rats (7 to 8-week-old, 200 to 300 g) and fixed using a dental resin. A tip of the guide cannula was positioned 0.9 mm behind a bregma, 1.2 mm at the right of a median line and in the depth of 1.5 mm from the brain surface. An inner needle was inserted such that its tip projected from the tip of the guide cannula by about 2 mm and arrived to a lateral cerebral ventricle. After a recovery period of about one week, a bovine pancreatic polypeptide (bPP, 5 μg/head/10μl) was administered to the lateral cerebral ventricle. A compound to be tested was simultaneously administered as a mixture with bPP. Food intake during 2 hours from the administration was measured. In this connection, both bPP and the compound to be tested were administered after dissolving them in 50 % propylene glycol.

[0104] The compound of the present invention significantly inhibits the increase in food intake induced by bPP (a homologue of NPY) simultaneously administered.

[0105] In consequence, the compound [I] of the present invention is useful as an agent for the treatment of various diseases associated with NPY, for example, diseases of circulatory organs such as hypertension, nephropathy, cardiopathy and angiospasm, diseases of central nervous system such as bulimia, depression, epilepsy and dementia, metabolic diseases such as obesity, diabetes and dysendocrisis, or glaucoma, especially bulimia, obesity and diabetes.

[0106] The compound represented by the general formula [I] can be administered orally or parenterally and by formulating into any dosage forms suitable for such an administration, it can be used as an agent for the treatment of the diseases of circulatory organs such as hypertension, nephropathy, cardiopathy and angiospasm, the diseases of central nervous system such as bulimia, depression, epilepsy and dementia, the metabolic diseases such as obesity, diabetes and dysendocrisis, or glaucoma. In clinical use of the compound of the present invention, it is also possible to administer the compound after formulating it into various pharmaceutical preparations by adding any pharmaceutically acceptable additive(s). Examples of such additive include those which are generally used in the field of pharmaceutical preparations, such as gelatin, lactose, sucrose, titanium oxide, starch, crystalline cellulose, hydroxypropylmethyl cellulose, carboxymethyl cellulose, corn starch, microcrystalline wax, white soft paraffine, magnesium aluminate methasilicate, anhydrous calcium phosphate, citric acid, trisodium citrate, hydroxypropyl cellulose, sorbitol, sorbitan fatty acid ester, polysorbate, sucrose fatty acid ester, polyoxyethylene hydrogenated castor oil, polyvinyl pyrrolidone, magnesium stearate, light anhydrous silicic acid, talc, vegetable oil, benzyl alcohol, gum arabic, propylene glycol, polyalkylene glycol, cyclodextrin and hydroxypropyl cyclodextrin.

[0107] Examples of the dosage form to be formulated as a mixture with these additives include solid preparations such as tablet, capsule, granule, powder or suppository; and liquid preparations such as syrup, elixir or injection, which can be prepared in accordance with any conventional method in the field of pharmaceuticals. In this connection, in the case of a liquid preparation, it may be in a form which is dissolved or suspended in water or other suitable solvent when used. Also, particularly in the case of an injection, it may be dissolved or suspended in physiological saline or glucose solution if necessary or further mixed with buffer and/or preservative.

[0108] The pharmaceutical preparation may contain the compound of the present invention in an amount of from 1.0 to 100% by weight, preferably from 1.0 to 60% by weight, with respect to the total preparation. The pharmaceutical preparation may also contain any other therapeutically effective compounds.

[0109] When the compound of the present invention is, for example, clinically used, its dosage and the number of times of its administration vary depending on the sex, age, body weight and the conditions of each patient and the nature and ranges of the intended therapeutic effects and the like. When it is administered to an adult, it is desirable in general to orally administer in an amount of from 0.1 to 100 mg/kg per day by dividing the daily dose into 1 to several times per day, or to parenterally administer in an amount of from 0.001 to 10 mg/kg by dividing the daily dose into 1 to several times per day.

BEST MODE FOR CARRYING OUT THE INVENTION

[0110] The present invention is described further in detail with reference to the following examples, but the invention should in no way be restricted thereby.

Example 1

Preparation of 5-(4-methoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

(1) Preparation of 5-(4-methoxyphenyl)-3-phenoxy carbonylaminopyrazole

[0111] Phenyl chlorocarbonate (0.87 ml) was added to a solution of 3-amino-5-(4-methoxyphenyl)pyrazole (1.19 g) in pyridine (10 ml), and the mixture was stirred at room temperature for 6 hours. The reaction solution was partitioned between ethyl acetate (50 ml) and water (30 ml), and the organic layer was washed with 1 N hydrochloric acid (50 ml x 2), saturated aqueous sodium bicarbonate (50 ml) and saturated brine (50 ml) and then dried over anhydrous magnesium sulfate. Then, the solvent was distilled away. The resulting residue was crystallized from ethyl acetate/hexane to obtain the title compound (0.95 g) as pale yellow crystals (melting point: 286 - 291°C).

(2) Preparation of 5-(4-methoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

[0112] A mixture of 5-(4-methoxyphenyl)-3-phenoxy carbonylaminopyrazole (93 mg), 1-phenylpiperazine (54 mg) and triethylamine (91 mg) was heated under reflux for 2 hours in chloroform (3 ml). After the reaction solution was allowed to cool, ethyl ether (3 ml) was added thereto, and the thus precipitated crystals were collected by filtration and dried to obtain the title compound (103 mg) as colorless crystals (melting point: 235 - 245°C).

[0113] Each of compounds of Examples 2 to 32 was obtained in the same manner as that described in Example 1 except that the starting materials used in Example 1 were replaced with appropriate starting materials corresponding to the desired compound.

Example 2

5-(4-chlorophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

[0114]

melting point: 245 - 248°C

Example 3

5-(2-methylphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

[0115]

melting point: 123 - 127°C

Example 4

5-(3-methylphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

[0116]

melting point: 140 - 141°C

Example 5

5-(4-methylphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

5 [0117]

melting point: 233 - 235°C

Example 6

10

5-(2-methoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

[0118]

15 melting point: 183 - 185°C

Example 7

5-(3,4-dichlorophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

20

[0119]

melting point: 252 - 254°C

25 Example 8

5-(4-bromophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

[0120]

30

melting point: 250 - 252°C

Example 9

35 5-(3-chlorophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

[0121]

melting point: 226 - 228°C

40

Example 10

5-phenyl-3-(4-phenylpiperazinyl)carbonylaminopyrazole

45 [0122]

melting point: 232°C

Example 11

50

5-(4-dimethylaminophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

[0123]

55 melting point: 242 - 246°C

Example 12

5-(3-dimethylaminophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

5 [0124]

melting point: 155 - 163°C (dihydrochloride)

Example 13

10

5-(3,4-dimethoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

[0125]

15 melting point: 116 - 119°C

Example 14

5-(4-isopropoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

20

[0126]

melting point: 245 - 246°C

25 Example 15

5-(4-ethoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

[0127]

30

melting point: 231 - 235°C

Example 16

35 5-(3-bromophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

[0128]

melting point: 208 - 209°C

40

Example 17

3-[4-(3-chlorophenyl)piperazinyl]carbonylamino-5-(4-methoxyphenyl)pyrazole

45 [0129]

melting point: 214 - 216°C

Example 18

50

3-[4-(3-chlorophenyl)piperazinyl]carbonylamino-5-(3,4-dimethoxyphenyl)pyrazole

[0130]

55 melting point: 115 - 117°C

Example 19

5-(2-chlorophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

5 [0131]

melting point: 168 - 169°C

Example 20

10

5-(4-methylthiophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

[0132]

15 melting point: 253 - 258°C

Example 21

5-(3,4-methylenedioxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

20

[0133]

melting point: 223 - 228°C

25 Example 22

3-(4-phenylpiperazinyl)carbonylamino-5-(4-pyridyl)pyrazole

[0134]

30

melting point: 249 - 255°C

Example 23

35 5-(4-methoxyphenyl)-3-methyl[2-(methylphenylamino)ethyl]aminocarbonylaminopyrazole

[0135]

40 ¹H-NMR(CDCI₃) δ: 2.99 (3H, s), 3.01 (3H, s), 3.56 (4H, s), 3.83 (3H, s), 6.23 (1H, brs), 6.75 - 6.85 (3H, m), 6.92 (2H, d, J=8.8Hz), 7.22 - 7.32 (2H, m), 7.45 (1H, brs), 7.54 (2H, d, J=8.8 Hz)

Example 24

5-(3-methoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

45

[0136]

melting point: 172 - 174°C

50 Example 25

5-(4-methoxyphenyl)-3-(2-methyl-4-phenylpiperazinyl)carbonylaminopyrazole

[0137]

55

melting point: 111 - 116°C

Example 26

5-(4-methoxyphenyl)-3-[2-(phenylamino)ethyl]aminocarbonylaminopyrazole

5 [0138]

melting point: 183 - 184°C

Example 27

10

5-(4-biphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

[0139]

15 melting point: 274 - 276°C

Example 28

20

5-(3-dimethylamino-4-methoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole

[0140]

melting point: 233 - 235°C

25 Example 29

3-(4-hydroxy-4-phenylpiperidino)carbonylamino-5-(4-methoxyphenyl)pyrazole

[0141]

30

melting point: 234 - 236°C

Example 30

35

5-(4-methoxyphenyl)-3-(4-phenyl-1,2,3,6-tetrahydropyridin-1-yl)carbonylaminopyrazole

[0142]

melting point: 232 - 235°C

40

Example 31

5-(4-methoxyphenyl)-3-(4-phenylpiperidino)carbonylaminopyrazole

45 [0143]

melting point: 205 - 208°C

Example 32

50

5-(4-methoxyphenyl)-3-(4-phenylhomopiperazinyl)carbonylaminopyrazole


[0144]

55 melting point: 115 - 123°C


THIS PAGE BLANK (USPTO)

5-(4-methoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-chlorophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(2-methylphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(3-methylphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-methylphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(2-methoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(3,4-dichlorophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-bromophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(3-chlorophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-phenyl-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-dimethylaminophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(3-dimethylaminophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(3,4-dimethoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-isopropoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-ethoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(3-bromophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 3-(4-(3-chlorophenyl)piperazinyl)carbonylamino-5-(4-methoxyphenyl)pyrazole,
 3-(4-(3-chlorophenyl)piperazinyl)carbonylamino-5-(3,4-dimethoxyphenyl)pyrazole,
 5-(2-chlorophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-methylthiophenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(3,4-methylenedioxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 3-(4-phenylpiperazinyl)carbonylamino-5-(4-pyridyl)pyrazole,
 5-(4-methoxyphenyl)-3-methyl[2-(methylphenylamino)ethyl]-aminocarbonylaminopyrazole,
 5-(3-methoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-methoxyphenyl)-3-(2-methyl-4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(4-methoxyphenyl)-3-[2-(phenylamino)ethyl]aminocarbonylaminopyrazole,
 5-(4-biphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 5-(3-dimethylamino-4-methoxyphenyl)-3-(4-phenylpiperazinyl)carbonylaminopyrazole,
 3-(4-hydroxy-4-phenylpiperidino)carbonylamino-5-(4-methoxyphenyl)pyrazole,
 5-(4-methoxyphenyl)-3-(4-phenyl-1,2,3,6-tetrahydropyridin-1-yl)carbonylaminopyrazole,
 5-(4-methoxyphenyl)-3-(4-phenylpiperidino)carbonylaminopyrazole, or
 5-(4-methoxyphenyl)-3-(4-phenylhomopiperazinyl)carbonylaminopyrazole.

10. A method for preparing a compound represented by the general formula [I]:




wherein A represents a nitrogen atom or a group represented by C-R⁵; Ar¹ represents an aryl group which may have a substituent selected from the group consisting of a halogen atom and lower alkyl and lower haloalkyl groups; Ar² represents an aryl or heteroaryl group which may have a substituent selected from the group consisting of a halogen atom and lower alkyl, lower alkenyl, lower haloalkyl, lower alkoxy, lower alkylthio, lower alkylamino, lower dialkylamino and aryl groups; R¹ represents a hydrogen atom, a lower alkyl group or a bond formed by linking to R⁵; R² represents a hydrogen atom or a lower alkyl group; R³ and R⁴ may be the same or different and each represents a hydrogen atom or a lower alkyl group, or R³ and R⁴ are linked to each other to form an alkylene group containing 2 to 4 carbon atoms which may have a lower alkyl group; and R⁵ represents a hydrogen atom or a hydroxyl, lower alkyl or lower alkoxy group or a bond formed by linking to R¹, which comprises reacting a compound represented by the general formula [II]:




[II]

wherein Ar^2 is as defined above,
with a compound represented by the general formula [III]:




[III]

wherein Ar^3 represents a phenyl group which may be substituted with a halogen atom or a nitro group; and X represents a halogen atom, to obtain a compound represented by the general formula [IV]:



[IV]


wherein Ar^2 and Ar^3 are as defined above,
and then reacting the compound [IV] with a compound represented by the general formula [V]:



[V]

wherein A, Ar^1 , R^1 , R^2 , R^3 and R^4 are as defined above.

11. A neuropeptide Y receptor antagonist comprising a compound represented by the general formula [I]:




[I]

wherein A represents a nitrogen atom or a group represented by $\text{C}-\text{R}^5$; Ar^1 represents an aryl group which may have a substituent selected from the group consisting of a halogen atom and lower alkyl and lower haloalkyl groups; Ar^2 represents an aryl or heteroaryl group which may have a substituent selected from the group consisting of a

halogen atom and lower alkyl, lower alkenyl, lower haloalkyl, lower alkoxy, lower alkylthio, lower alkylamino, lower dialkylamino and aryl groups; R^1 represents a hydrogen atom, a lower alkyl group or a bond formed by linking to R^5 ; R^2 represents a hydrogen atom or a lower alkyl group; R^3 and R^4 may be the same or different and each represents a hydrogen atom or a lower alkyl group, or R^3 and R^4 are linked to each other to form an alkylene group containing 2 to 4 carbon atoms which may have a lower alkyl group; and R^5 represents a hydrogen atom or a hydroxyl, lower alkyl or lower alkoxy group or a bond formed by linking to R^1 , or a salt thereof as an active ingredient.

12. An agent for the treatment of bulimia, obesity or diabetes comprising a compound represented by the general formula [I]:



wherein A represents a nitrogen atom or a group represented by $C-R^5$; Ar^1 represents an aryl group which may have a substituent selected from the group consisting of a halogen atom and lower alkyl and lower haloalkyl groups; Ar^2 represents an aryl or heteroaryl group which may have a substituent selected from the group consisting of a halogen atom and lower alkyl, lower alkenyl, lower haloalkyl, lower alkoxy, lower alkylthio, lower alkylamino, lower dialkylamino and aryl groups; R^1 represents a hydrogen atom, a lower alkyl group or a bond formed by linking to R^5 ; R^2 represents a hydrogen atom or a lower alkyl group; R^3 and R^4 may be the same or different and each represents a hydrogen atom or a lower alkyl group, or R^3 and R^4 are linked to each other to form an alkylene group containing 2 to 4 carbon atoms which may have a lower alkyl group; and R^5 represents a hydrogen atom or a hydroxyl, lower alkyl or lower alkoxy group or a bond by linking to R^1 , or a salt thereof as an active ingredient.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP97/04399

A. CLASSIFICATION OF SUBJECT MATTER Int.Cl. ⁶ C07D231/40, 401/01, 401/12, 405/12, A61K31/415, 31/445, 31/495, 31/55 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int.Cl. ⁶ C07D231/40, 401/01, 401/12, 405/12, A61K31/415, 31/445, 31/495, 31/55 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CA (STN), REGISTRY (STN)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JP, 48-10069, A (Takeda Chemical Industries, Ltd.), February 8, 1973 (08. 02. 73) (Family: none)	1 - 12
A	JP, 7-188269, A (Otsuka Pharmaceutical Factory, Inc.), July 25, 1995 (25. 07. 95) (Family: none)	1 - 12
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search January 29, 1998 (29. 01. 98)		Date of mailing of the international search report February 10, 1998 (10. 02. 98)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.